

Data reported as: % susceptible (# isolates tested)¹

Antibiotic	B bron	E coli	E fael	E faem	Ente	K pneu	P aer	P mult	Pseu	S aur	S can	S pint
Amikacin	100% (22)	99% (711)	23% (227)	13% (40)	100% (99)	96% (54)	92% (268)	97% (35)	85% (102)	100% (38)	3% (254)	100% (869)
Amoxicillin/Clavulanic Acid	91% (22)	84% (711)	97% (227)	33% (40)	33% (99)	74% (54)	1% (268)	100% (35)	47% (102)	82% (38)	100% (254)	85% (869)
Ampicillin	14% (22)	71% (711)	97% (227)	33% (40)	35% (99)	13% (54)	1% (268)	100% (35)	42% (102)	26% (38)	100% (254)	46% (869)
Cefazolin	0% (22)	86% (711)	2% (227)	8% (40)	26% (99)	74% (54)	2% (268)	100% (35)	34% (102)	82% (38)	100% (254)	85% (869)
Cefovecin	0% (22)	86% (711)	1% (227)	5% (40)	81% (99)	81% (54)	1% (268)	100% (35)	37% (102)	82% (38)	98% (254)	81% (869)
Cefoxitin	0% (22)	88% (711)	0% (227)	3% (40)	37% (99)	78% (54)	1% (268)	100% (35)	40% (102)	55% (38)	100% (254)	84% (869)
Cefpodoxime	0% (22)	86% (711)	11% (227)	8% (40)	85% (99)	83% (54)	1% (268)	97% (35)	35% (102)	74% (38)	99% (254)	82% (869)
Ceftiofur	0% (22)	87% (711)	5% (227)	8% (40)	86% (99)	83% (54)	2% (268)	100% (35)	35% (102)	82% (38)	99% (254)	83% (869)
Cephalothin	50% (2)	100% (4)	3% (68)	30% (10)	0% (1)		0% (1)	100% (1)	50% (2)	82% (38)	99% (246)	84% (825)
Chloramphenicol	95% (22)	87% (711)	93% (227)	90% (40)	85% (99)	81% (54)	2% (268)	100% (35)	53% (102)	79% (38)	98% (254)	87% (869)
Clindamycin	0% (22)	0% (711)	0% (227)	25% (40)	0% (99)	0% (54)	0% (268)	3% (35)	13% (102)	92% (38)	89% (254)	74% (869)
Doxycycline	100% (22)	84% (711)	73% (227)	50% (40)	80% (99)	76% (54)	6% (268)	97% (35)	75% (102)	89% (38)	65% (254)	60% (869)
Enrofloxacin	95% (22)	91% (711)	31% (227)	3% (40)	95% (99)	85% (54)	46% (268)	97% (35)	67% (102)	87% (38)	48% (254)	79% (869)
Erythromycin	0% (22)	0% (711)	32% (227)	5% (40)	1% (99)	0% (54)	0% (268)	29% (35)	25% (102)	61% (38)	0% (254)	73% (869)
Gentamicin	82% (22)	93% (711)	44% (227)	5% (40)	99% (99)	91% (54)	76% (268)	100% (35)	83% (102)	100% (38)	42% (254)	81% (869)
Imipenem	100% (22)	100% (711)	98% (227)	30% (40)	99% (99)	100% (54)	97% (268)	100% (35)	98% (102)	79% (38)	100% (254)	85% (869)
Marbofloxacin	95% (22)	92% (711)	32% (227)	3% (40)	96% (99)	93% (54)	77% (268)	100% (35)	87% (102)	87% (38)	77% (254)	84% (869)
Oxacillin ³	NI	NI	NI	NI	NI	NI	NI	NI	NI	82% (38)	NI	81% (869)
Penicillin	0% (22)	0% (711)	98% (227)	28% (40)	0% (99)	0% (54)	0% (268)	74% (35)	1% (102)	21% (38)	98% (254)	31% (869)
Ticarcillin	55% (22)	72% (711)	7% (227)	13% (40)	72% (99)	13% (54)	92% (268)	100% (35)	65% (102)	79% (38)	100% (254)	84% (869)
Ticarcillin/Clavulanic Acid	95% (22)	84% (711)	5% (227)	13% (40)	89% (99)	81% (54)	93% (268)	100% (35)	72% (102)	82% (38)	99% (254)	84% (869)
Trimethoprim/Sulphamethoxazole	32% (22)	89% (711)	89% (227)	80% (40)	94% (99)	83% (54)	9% (268)	97% (35)	48% (102)	100% (38)	98% (254)	78% (869)

³ Isolates resistant to oxacillin are interpreted as potentially methicillin resistant.

Key:

1	Data is reported as: % susceptible (# isolates tested) - not all bacteria isolated at ISU VDL have been tested for antimicrobial susceptibility	
2	See Salmonella serotype table for most common serotypes isolated within each group	
3	Isolates resistant to oxacillin are interpreted as potentially methicillin resistant.	
4	A result of ≤ 2 ug/ml for Carbadox is a conservative indicator of bacterial inhibition by this antimicrobial agent. The result shown is based on pharmacokinetic research indicating an average Carbadox level of 4.5 mcg/ml in the small intestine of pigs fed a dose rate of 50 g/ton. (De Graff 1988).	
5	Multidrug resistant isolates were found resistant to most classes of antimicrobial in the 1 st round of testing. This table represents additional Disk Diffusion testing for those isolates.	
NA	Not applicable	
ND	Not done	
NI	No interpretation	
A equ - Actinobacillus equuli	H ecol - hemolytic E.coli	S aur - Staphylococcus aureus
A suis - Actinobacillus suis	H som - Histophilus somni	S beta- Beta Streptococcus species
Abua - Acinetobacter species	HPS - Haemophilus parasuis	S can - Streptococcus canis
Amy - Actinomyces species	K pneu - Klebsiella pneumoniae	S chol - Salmonella choleraesuis
APP - Actinobacillus pleuropneumoniae	M bov - Moraxella bovis	S dysg - Streptococcus dysgalactiae
B bron - Bordetella bronchiseptica	M haem - Mannheimia haemolytica	S epi- Staphylococcus epidermidis
B tre - Bibersteinia trehalosi (formerly Pasteurella trehalosi)	P aer - Pseudomonas aeruginosa	S equi - Streptococcus equi
Bact - Bacteroides group	P cab - Pasteurella caballi	S equus - Streptococcus equisimilis
C diff - Clostridium difficile	P mult - Pasteurella multocida	S pint - Staph pseudintermedius
C perf - Clostridium perfringens	Past - Pasteurella species	S suis - Streptococcus suis
Clos - Clostridium species	Pec - Peptococcus species	S ube - Streptococcus uberis
E coli - Escherichia coli	Pes - Peptostreptococcus species	S zoo - Streptococcus zooepidemicus
E fael - Enterococcus faecalis	Pmul A - Pasteurella multocida Type A	Salm sp- Salmonella species
E faem - Enterococcus faecium	Pmul D - Pasteurella multocida Type D	Salm B - Salmonella species group B
Enc - Enterococcus species	Prot - Proteus species	Salm C1 - Salmonella species group C1
Ente - Enterobacter species	Prp - Propionibacterium species	Salm C2 - Salmonella species group C2
Erys - Erysipelothrix	Pseu - Pseudomonas species	Salm D - Salmonella species group D
Fus - Fusobacterium	R equ - Rhodococcus equi	Salm E - Salmonella species group E
G ana - Gallibacterium anatis		