### We're Here to Help

The Iowa State University
Veterinary Diagnostic
Laboratory has available an
immunohistochemical
test for BVDV persistent
infection on formalin-fixed ear
notch samples.

If any questions arise regarding this or any testing at ISU-VDL, please feel free to contact one of our diagnosticians.



# Immunohistochemistry for BVDV Persistent Infection



IOWA STATE UNIVERSITY
VETERINARY DIAGNOSTIC LABORATORY



IOWA STATE UNIVERSITY
VETERINARY DIAGNOSTIC LABORATORY
1600 S 16th Street
College of Veterinary Medicine
Iowa State University
Ames, IA 50011

515-294-1950 www.vdpam.iastate.edu

lowa State University does not discriminate on the basis of race, color, age, religion, national origin, sexual orientation, gender identity, sex, marital status, disability, or status as a U.S. veteran. Inquiries can be directed to the Director of Equal Opportunity and Diversity, 3680 Beardshear Hall, (515) 294-7612.

## BVDV Surveillance Using Ear Notch Immunohistochemistry

Immunohistochemistry on ear notch samples pinpoints BVDV persistent infection. It is appropriate for cattle of all ages and is considered the test of choice for young animals.

Suggestions from the Academy of Veterinary Consultants have been published to help practitioners and producers in making decisions related to BVDV control strategies for their herds.

Immunohistochemistry on ear notch samples is one potential testing option.



### Why Test?

Good candidates for BVDV surveillance testing are herds which exhibit the following:

Poor reproductive performance

High calf morbidity/mortality despite good sanitation and nutrition

Laboratory testing confirming evidence of BVDV in the herd

For optimum results, testing should start BEFORE the breeding season, producer goals should be known and achievable, and a plan of action should be in place.

#### Who Needs to be Tested?

In general:

- All calves should be tested
- All open cows or cows which have lost calves
- All replacement bulls and cows
- All replacement heifers



Please bear in mind that negative BVDV Pl testing does not guarantee absence of a BVDV problem.

Without adequate attention to vaccination programs and biosecurity concerns, simply removing PI cattle from the herd will not necessarily remove a BVDV or any other disease problem.

Feel free to contact Dr. Vickie Cooper or any of the diagnostic pathologists at ISU-VDL for help in tailoring a herd health surveillance program with emphasis on BVDV for your producers.

